

Power Law Distributions of Seismic Rates

A. Saichev^{1,2} and D. Sornette^{3,4}

¹*Mathematical Department, Nizhny Novgorod State University,
Gagarin prosp. 23, Nizhny Novgorod, 603950, Russia*

²*Institute of Geophysics and Planetary Physics, University of California, Los Angeles, CA 90095*

³*Institute of Geophysics and Planetary Physics and Department of Earth and Space Sciences,
University of California, Los Angeles, CA 90095*

⁴*Laboratoire de Physique de la Matière Condensée,*

*CNRS UMR 6622 and Université de Nice-Sophia Antipolis, 06108 Nice Cedex 2, France**

(Dated: February 2, 2008)

We report an empirical determination of the probability density functions $P_{\text{data}}(r)$ of the number r of earthquakes in finite space-time windows for the California catalog. We find a stable power law tail $P_{\text{data}}(r) \sim 1/r^{1+\mu}$ with exponent $\mu \approx 1.6$ for all space (5×5 to 20×20 km²) and time intervals (0.1 to 1000 days). These observations, as well as the non-universal dependence on space-time windows for all different space-time windows simultaneously, are explained by solving one of the most used reference model in seismology (ETAS), which assumes that each earthquake can trigger other earthquakes. The data imposes that active seismic regions are Cauchy-like fractals, whose exponent $\delta = 0.1 \pm 0.1$ is well-constrained by the seismic rate data.

PACS numbers: 91.30.Px ; 89.75.Da; 05.40.-a

Seismicity is perhaps the best example of a self-organizing process exhibiting scaling diagnosed with so many power laws: the Gutenberg-Richter distribution $\sim 1/E^{1+\beta}$ (with $\beta \approx 2/3$) of earthquake energies E ; the Omori law $\sim 1/t^p$ (with $p \approx 1$ for large earthquakes) of the rate of aftershocks as a function of time t since a mainshock; the productivity law $\sim E^a$ (with $a \approx 2/3$) giving the number of earthquakes triggered by an event of energy E [1]; the power law distribution $\sim 1/L^2$ of fault lengths L [2]; the fractal structure of fault networks [3] and of the spatial organization of earthquake epicenters [4]; the distribution $1/s^{2+\delta}$ (with $\delta \geq 0$) of seismic stress sources s in earthquake focal zones due to past earthquakes [5]. Related universal laws for the distribution of waiting times and seismic rates between earthquakes have recently been derived from the analyses of space-time windows [6].

Here, we report and explain theoretically an addition empirical power law: the numbers r of earthquakes in finite space-time windows for the California SCEC catalog, over fixed spatial boxes 5×5 km² to 20×20 km² and time intervals $dt = 1, 10, 100$ and 1000 days are distributed with a stable power law tail $P_{\text{data}}(r) \sim 1/r^{1+\mu}$ with exponent $\mu \approx 1.6$ for all time intervals. This result has important implications in constraining the physics of earthquakes and in estimating the performance of forecasting models of seismicity. For the former, we show that this observation can be rationalized by a simple stochastic branching model (ETAS) which uses the above mentioned Gutenberg-Richter, Omori, and productivity laws applied to a fractal spatial geometry of earthquake epicenters. The fundamental physical ingredient is that each earthquake can trigger other earthquakes (“aftershocks”) and an earthquake sequence results in this model from the cascade of aftershocks triggered by each past earth-

quake. In addition, the growing efforts in earthquake forecasts requires estimating the performance of a forecasting model by a likelihood function, which are currently based on Poisson probabilities calculated using declustered catalogues. Our work shows that spontaneous fluctuations of the number of triggered earthquakes in space-time bins, due to the cascades of triggering processes, may lead to dramatic departures from the Poisson model used as one of the building block of standard testing procedures.




FIG. 1: Empirical probability density functions (pdf) $P_{\text{data}}(r)$ of the number r of earthquakes in the space-time bins of size 5×5 km² and $dt = 1$ days of the SCEC catalog. The straight line is the best fit with a pure power law (1). The continuous line is the theoretical prediction derived below for $dt = 1$ days and with no adjustable parameters, as the parameters are fixed to the values adjusted from the fit for $dt = 1000$ days for which the pdf is strongly curved.

In order to maximize the size and quality of the data used for our analysis, we consider the time interval 1994 – 2003 in a region from approximately 32° to 37° N in latitude, and from -114° to -122° in longitude, of the Southern Californian earthquakes catalog with revised magnitudes $M_L > 1.5$, which contains a total of 86,228 earthquakes. The spatial domain is covered by square boxes of $(L = 5 \text{ km}) \times (L = 5 \text{ km})$. Other larger box sizes given similar results. We consider time windows from $dt = 1$ day to $dt = 1000$ days. Figure 1 plots the empirical probability density functions $P_{\text{data}}(r)$ of the number r of earthquakes in the space-time bins described above for $dt = 1$ days. The straight line is the best fit with a pure power law

$$P_{\text{data}}(r) \sim 1/r^{1+\mu} \quad (1)$$

over the range $1 \leq r \leq 100$. Similar power law tails are present in the tail for the other time windows. The estimation for μ is stable since the fitted values are $\mu = 1.65$ for $dt = 100$ days, $\mu = 1.75$ for $dt = 10$ days and $\mu = 1.60$ for $dt = 1$ days. However, the pdf becomes more and more curved in the larger portion of the bulk as the size dt of the time window is increased [7]. This behavior can be explained by the theory described below.

The ETAS (Epidemic-Type Aftershock Sequence) model of triggered seismicity is based on the three first well-founded empirical laws mentioned above. Its appeal lies in its simplicity, its power of explanation of other empirical observations (see for instance [8] and references therein) and its wide use as a benchmark. The ETAS model belongs to the general class of branching processes with infinite variance of the number of progenies per mother, with a long-time (power law) memory of the impact of a mother on her first-generation daughters described by the empirical Omori law for aftershocks. These two ingredients together with the mechanism of cascades of branching have been shown to give rise to subdiffusion and to non mean-field behavior in the distribution of the total number of aftershocks per mainshock [9], in the distribution of the total number of generations before extinctions and in the distribution of the total duration of an aftershock sequence before extinction [10].

In the ETAS model, each earthquake is a potential progenitor or mother, characterized by its conditional average number $N_m \equiv \kappa\mu(m)$ of children (triggered events or aftershocks of first generation), where $\mu(m) = 10^{\alpha(m-m_0)}$ is proportional to the average productivity of an earthquake of magnitude $m \geq m_0$ [1], κ is a constant factor and m_0 is the minimum magnitude of earthquakes capable of triggering other earthquakes. For a given earthquake of magnitude m and therefore of mark $\mu(m)$, the number r of its daughters of first generation are drawn at random according to the Poissonian statistics $p_\mu(r) = \frac{N_m^r}{r!} e^{-N_m} = \frac{(\kappa\mu)^r}{r!} e^{-\kappa\mu}$. The challenge of our present analysis is to understand how the exponential Poisson distribution is renormalized into the power law

(1) by taking into account all earthquake triggering paths simultaneously over all possible generations. The ETAS model is complemented by the normalized Gutenberg-Richter (GR) density distribution of earthquake magnitudes $p(m) = b \ln(10) 10^{-b(m-m_0)}$, $m \geq m_0$. This magnitude distribution $p(m)$ is assumed to be independent of the magnitude of the triggering earthquake, i.e., a large earthquake can be triggered by a smaller one. Combining the GR and the productivity laws shows that the earthquake marks μ and therefore the conditional average number N_m of daughters of first generation are distributed according to the normalized power law

$$p_\mu(\mu) = \frac{\gamma}{\mu^{1+\gamma}}, \quad 1 \leq \mu < +\infty, \quad \gamma = b/\alpha. \quad (2)$$

For earthquakes, $b \approx 1$ and $0.5 < \alpha < 1$ giving $1 < \gamma < 2$. This range $1 < \gamma < 2$ implies that the mathematical expectation of μ and therefore of N_m (performed over all possible magnitudes) is finite but its variance is infinite. Given γ , the coefficient κ then controls the value of the average number n (or branching ratio) of children of first generation per mother: $n = \langle N_m \rangle = \kappa \langle \mu \rangle = \kappa \frac{\gamma}{\gamma-1}$, where the average $\langle N_m \rangle$ is taken over all mothers' magnitudes drawn from the GR law. Recall that the values $n < 1$, $n = 1$ and $n > 1$ correspond respectively to the sub-critical, critical and super-critical branching regimes. The last ingredient of the ETAS model consists in the specification of the space-time rate function $N_m \Phi(\mathbf{r} - \mathbf{r}_i, t - t_i)$ giving the average rate of first generation daughters at time t and position $\mathbf{r}$ created by a mother of magnitude $m \geq m_0$ occurring at time t_i and position $\mathbf{r}_i$. We use the standard factorization $\Phi(\mathbf{x}, t) = \Phi(t) \phi(\mathbf{x})$. The time propagator $\Phi(t)$ has the Omori law form $\Phi(t) = \frac{\theta c^\theta}{(c+t)^{1+\theta}} H(t)$ where $H(t)$ is the Heaviside function, $0 < \theta < 1$, c is a regularizing time scale that ensures that the seismicity rate remains finite close to the mainshock. The space propagator is $\phi(\mathbf{x}) = \frac{\eta d^\eta}{2\pi(x^2+d^2)^{(\eta+2)/2}}$. The next ingredient of the ETAS model is to assume that plate tectonic motion induces spontaneous mother earthquakes, which are not triggered by previous earthquakes, according to a Poissonian point process, such that the average number of spontaneous mother earthquakes per unit time and per unit surface is ϱ . In the ETAS branching model, each such spontaneous mother earthquake then triggers independently its own space-time aftershocks branching process. The last ingredient of our theory is to recognize that, at large scale, earthquakes are preferentially clustered near the plate boundaries while, at smaller scales, earthquakes are found mostly along faults and close to nodes between several faults. We thus extend slightly the ETAS model to allow for the heterogeneity of the spontaneous earthquake sources ϱ reflecting the influence of pre-existing fault structures, some rheological heterogeneity and complex spatial stress distributions. For this, we use the distribution of the stress field in heteroge-

neous media and due to earthquakes [5] which is found close to a Cauchy distribution. The simplest prescription is to assume that ϱ is itself random and distributed according to $\frac{1}{\langle \varrho \rangle} f\left(\frac{\varrho}{\langle \varrho \rangle}\right)$, where $\langle \varrho \rangle$ is then statistical average of the random space-time Poissonian source intensity ϱ . In the numerical applications, we shall use the form $f_\delta(x) = \frac{\delta+1}{\delta} \left(1 + \frac{x}{\delta}\right)^{-1-\delta}$ ($\delta > 0$).

Due to the independence between each sequence triggered by each spontaneous event, the generating probability function (GPF) of the number of events (including mother earthquakes and all their aftershocks of all generations), falling into the space-time window $\{[t, t+\tau] \times \mathcal{S}\}$ is equal to

$$\Theta_{\text{sp}}(z, \tau, \mathcal{S}) = e^{-\varrho L(z, \tau, \mathcal{S})} \quad (3)$$

where $L(z, \tau, \mathcal{S}) = \int_0^\infty dt \iint_{-\infty}^\infty d\mathbf{x} [1 - \Theta(z, t, \tau, \mathcal{S}; \mathbf{x})] + \int_0^\tau dt \iint_{-\infty}^\infty d\mathbf{x} [1 - \Theta(z, t, \mathcal{S}; \mathbf{x})][1 - I_{\mathcal{S}}(\mathbf{x})] + \int_0^\tau dt \iint_{\mathcal{S}} d\mathbf{x} [1 - z\Theta(z, t, \mathcal{S}; \mathbf{x})]$. The first summand in $L(z, \tau, \mathcal{S})$ describes the contribution to the GPF Θ_{sp} from aftershocks triggered by mother earthquakes that occurred before the time window (i.e. at instants t' such that $t' < t$). The corresponding GPF $\Theta(z, t - t', \tau, \mathcal{S}; \mathbf{x})$ of the number of aftershocks triggered inside the space-time window $\{[t, t+\tau] \times \mathcal{S}\}$ by some mother event that occurred at time $t' = 0$ satisfies the relation

$$\Theta(z, t, \tau, \mathcal{S}; \mathbf{x}) = G[1 - \Psi(z, t, \tau, \mathcal{S}; \mathbf{x})], \quad (4)$$

where the auxiliary function $\Psi(z, t, \tau, \mathcal{S}; \mathbf{x})$, describing the space-time dissemination of aftershocks triggering by some mother event, is equal to

$$\begin{aligned} \Psi(z, t, \tau, \mathcal{S}; \mathbf{x}) &= \Phi(\mathbf{x}, t) \otimes [1 - \Theta(z, t, \tau, \mathcal{S}; \mathbf{x})] \\ &+ \Phi(\mathbf{x}, t + \tau) \otimes [1 - \Theta(z, \tau, \mathcal{S}; \mathbf{x})] + \\ &(1 - z)\Phi(\mathbf{x}, t + \tau) \otimes I_{\mathcal{S}}(\mathbf{x})\Theta(z, \tau, \mathcal{S}; \mathbf{x}). \end{aligned} \quad (5)$$

The function $I_{\mathcal{S}}(\mathbf{x})$ in (5) is the indicator of the space window $\mathcal{S}$ and $G(z)$ in (4) is the GPF of the number R_1 given by (2) of first generation aftershocks triggered by some mother earthquake, given by $G(z) = \gamma\kappa^\gamma(1-z)^\gamma \Gamma(-\gamma, \kappa(1-z))$. The last two summands of $L(z, \tau, \mathcal{S})$ describe the contribution of aftershocks triggered by earthquakes, occurring inside the time window (i.e., $t' \in [t, t+\tau]$). The second (resp. third) term corresponds to the subset spatially outside (resp. inside) the domain $\mathcal{S}$. These last two terms depend on the GPF $\Theta(z, \tau, \mathcal{S}; \mathbf{x}) = \Theta(z, t = 0, \tau, \mathcal{S}; \mathbf{x})$ of the numbers of aftershocks triggered till time τ inside the space window $\mathcal{S}$ by some mother event arising at the instant $t = 0$ and at the point $\mathbf{x}$. It follows from (4) and (5) that it satisfies the relations

$$\Theta(z, \tau, \mathcal{S}; \mathbf{x}) = G[1 - \Psi(z, \tau, \mathcal{S}; \mathbf{x})] \quad (6)$$

and

$$\begin{aligned} \Psi(z, \tau, \mathcal{S}; \mathbf{x}) &= \Phi(\mathbf{x}, \tau) \otimes [1 - \Theta(z, \tau, \mathcal{S}; \mathbf{x})] \\ &+ (1 - z)\Phi(\mathbf{x}, \tau) \otimes I_{\mathcal{S}}(\mathbf{x})\Theta(z, \tau, \mathcal{S}; \mathbf{x}). \end{aligned} \quad (7)$$

In addition, we shall need the GPF

$$\Theta(z, \mathcal{S}; \mathbf{x}) = \Theta(z, \tau = \infty, \mathcal{S}; \mathbf{x}) \quad (8)$$

of the total numbers of aftershocks triggered by some mother earthquake inside the area $\mathcal{S}$. As seen from (6) and (7), it satisfies the relations

$$\Theta(z, \mathcal{S}; \mathbf{x}) = G[1 - \Psi(z, \mathcal{S}; \mathbf{x})] \quad (9)$$

and $\Psi(z, \mathcal{S}; \mathbf{x}) = 1 - \phi(\mathbf{x}) \otimes \Theta(z, \mathcal{S}; \mathbf{x}) + (1 - z)\phi(\mathbf{x}) \otimes I_{\mathcal{S}}(\mathbf{x})\Theta(z, \mathcal{S}; \mathbf{x})$.

Taking into account the distribution of the source intensities ϱ amounts to averaging equation (3) over ϱ weighted with the statistics $\frac{1}{\langle \varrho \rangle} f\left(\frac{\varrho}{\langle \varrho \rangle}\right)$. This gives

$$\Theta_{\text{sp}}(z, \tau; \mathcal{S}) = \hat{f}[\langle \varrho \rangle L(z, \tau, \mathcal{S})], \quad (10)$$

where $\hat{f}(u)$ is the Laplace transform of the pdf $f(x)$.

To go further, we make two approximations. If the time duration τ of the space-time window is sufficiently large, it can be shown that the statistical averages of the seismic rates become independent of τ . It seems reasonable to conjecture that the GPF $\Theta(z, \tau, \mathcal{S}; \mathbf{x})$ of the total number of aftershocks triggered by some earthquake source inside the space domain $\mathcal{S}$ until time τ coincides approximately with the saturated GPF $\Theta(z, \mathcal{S}; \mathbf{x})$ of the total number of aftershocks triggered by some earthquake source inside the space domain $\mathcal{S}$. Within this approximation of large time windows, the effect of aftershocks triggered by earthquake sources occurring till the beginning t of the time window is negligible. Within this large time window approximation, one may ignore the first term in the contribution to (3) and replace $\Theta(z, t, \mathcal{S}; \mathbf{x})$ by $\Theta(z, \mathcal{S}; \mathbf{x})$ in the remaining terms. As a result, $L(z, \tau, \mathcal{S})$ in (3) takes the following approximate form $L(z, \tau, \mathcal{S}) \simeq \tau \iint_{-\infty}^\infty [1 - \Theta(z, \mathcal{S}; \mathbf{x})][1 - I_{\mathcal{S}}(\mathbf{x})]d\mathbf{x} + \tau \iint_{\mathcal{S}} [1 - z\Theta(z, \mathcal{S}; \mathbf{x})]d\mathbf{x}$, where $\Theta(z, \mathcal{S}; \mathbf{x})$ is the solution of $\Theta = G[\Theta \otimes \phi - (1 - z)I_{\mathcal{S}}\Theta \otimes \phi]$. To find a reasonable approximate expression for the sought GPF $\Theta(z, \mathcal{S}; \mathbf{x})$, notice that if the spatial extend ℓ of the window is larger than the characteristic scale d of the space kernel, or if n is close to 1, then the characteristic spatial scale associated with the GPF $\Theta(z, \mathcal{S}; \mathbf{x})$ becomes greater than d . Therefore, without essential error, one may replace $\Theta \otimes \phi$ by Θ . In addition, we take into account the finiteness of the domain $\mathcal{S}$ by using the factorization procedure: $I_{\mathcal{S}}(\mathbf{x})\Theta(z, \mathcal{S}; \mathbf{x}) \otimes \phi(\mathbf{x}) \simeq \Theta(z, \mathcal{S}; \mathbf{x}) p_{\mathcal{S}}(\mathbf{x})$, where $p_{\mathcal{S}}(\mathbf{x})$ remains to be specified. This amounts to replacing a convolution integral by an algebraic term. This factorization approximation is a crucial step of our analysis

and is justified elsewhere [7]. As a result of its use, the nonlinear integral equation for $\Theta(z, \mathcal{S}; \mathbf{x})$ transforms into the functional equation $\Theta = G[(1 + (z - 1)p_S(\mathbf{x}))\Theta]$. This approximation leads to $\langle R \rangle = \frac{n}{1-n} p_S$, where $\langle R \rangle$ is the average of the total number of events in the space-time window. The effective parameter p_S can be determined from the consistency condition such that $\langle R \rangle$ be equal to the true $\langle R(\mathcal{S}; \mathbf{x}) \rangle$, which can be calculated exactly. This gives

$$p_S(\mathbf{x}) = \frac{1-n}{n} \langle R(\mathcal{S}; \mathbf{x}) \rangle, \quad \tilde{p}_S(\mathbf{q}) = \tilde{I}_S(\mathbf{q}) \tilde{\phi}(\mathbf{q}) \frac{1-n}{1-n\tilde{\phi}(\mathbf{q})}. \quad (11)$$

For $\ell \gg d$, the factor $p_S(\mathbf{x})$ approaches a rectangular function. We can use this observation to help determine the statistics of the number of events in a finite space-time window, using the approximation $p_S(\mathbf{x}) \simeq \text{const} = p$ for $\mathbf{x} \in \mathcal{S}$. We define the parameter p as the space average of $p_S(\mathbf{x})$ over the window's area $\mathcal{S}$:

$$p \simeq \frac{1}{S} \iint_{\mathcal{S}} p_S(\mathbf{x}) d\mathbf{x}. \quad (12)$$

This approximation allows us to get

$$\tau \iint_{\mathcal{S}} [1 - z\Theta(z, \mathcal{S}; \mathbf{x})] d\mathbf{x} \simeq \tau S [1 - z\Theta(z; p)], \quad (13)$$

where $\Theta(z, p)$ is the solution of $\Theta(z; p) = G[(1 + (z - 1)p)\Theta(z; p)]$.

Complementarily, a study of $p_S(\mathbf{x})$ shows that it is small outside the window space domain $\mathcal{S}$. This implies that, outside $\mathcal{S}$, one may replace the functional equation on Θ by $1 - \Theta(z, \mathcal{S}; \mathbf{x}) \simeq \frac{n}{1-n} (1 - z)p_S(\mathbf{x})$. Therefore, we get $\tau \iint_{-\infty}^{\infty} [1 - \Theta(z, \mathcal{S}; \mathbf{x})] [1 - I_S(\mathbf{x})] d\mathbf{x} \simeq q\tau S \frac{n}{1-n} (1 - z)$,

where $q = \frac{1}{S} \iint_{-\infty}^{\infty} p_S(\mathbf{x}) [1 - I_S(\mathbf{x})] d\mathbf{x}$. Taking into account

that $\iint_{-\infty}^{\infty} p_S(\mathbf{x}) d\mathbf{x} = S$, we obtain $q \simeq 1 - p$. Putting all these approximations together allows us to rewrite the expression of $L(z, \tau, \mathcal{S})$ in (3) as

$$L(z, \tau, \mathcal{S}) \simeq \tau S \left[\frac{n}{1-n} (1 - p)(1 - z) + 1 - z\Theta(z; p) \right]. \quad (14)$$

The factorization procedure obtains the characteristic features of the space-time branching process in a finite space-time domain, at the cost of an adjustable parameter p .

Starting from the general expression (10) of the GPF $\Theta_{\text{sp}}(z, \tau; \mathcal{S})$ with the approximation (14) for $L(z, \tau, \mathcal{S})$ and using the relationship between the probability $P_{\text{sp}}(r; \rho, p)$ and its GPF in the form of its integral representation, we obtain the following expression valid in the limit of sufficiently large time windows $P_{\text{sp}}(r; \rho, p) =$

$$\frac{1}{2\pi i} \oint_{\mathcal{C}} \hat{f} \left(\rho \left[\frac{n}{1-n} (1 - p)(1 - z) + 1 - z\Theta(z; p) \right] \right) \frac{dz}{z^{r+1}}.$$

Introducing the new integration variable $y = (1 + (z - 1)p)\Theta(z; p) \iff z = Z(y) = \frac{1}{p} \left(\frac{y}{G(y)} + p - 1 \right)$, by construction of y , $\Theta(z; p) = G(y)$ which allows us to obtain the following explicit expression $P_{\text{sp}}(r; \rho, p) = \frac{1}{2\pi i} \times \oint_{\mathcal{C}'} \hat{f} \left(\rho \left[\frac{n}{1-n} (1 - p)(1 - Z(y)) + 1 - Z(y)G(y) \right] \right) \frac{dZ(y)}{dy} \frac{dy}{Z^{r+1}(y)}.$

This expression makes a precise quantitative prediction for the dependence of the distribution $P_{\text{sp}}(r; \rho, p)$ of the number r of earthquakes per space-time window as a function of r , once the following model parameters are given: the branching ratio n , the exponent γ of the distribution of productivities, the exponent δ of the distribution of spontaneous earthquake sources, the fraction p_S of direct (first generation) aftershocks which fall within the domain $\mathcal{S}$, and the average number ρ of spontaneous earthquake source per space-time bin defined by $\rho = \langle \varrho \rangle \tau S$. The theoretical curve in Fig. 1 is obtained by a numerical integration of $P_{\text{sp}}(r; \rho, p) =$ for the set of parameters $n = 0.96$, $\gamma = 1.1$, $p_S = 0.25$, $\delta = 0.15$ and $\rho = 0.0019 \text{ dt}$ with dt in units of days (thus equal to 100 for Fig. 1). These parameters give the best fit for the large time window $\text{dt} = 1000$ days. They have been kept fixed for the other time windows which exhibit very different shapes in their bulk. The theory is thus able to account simultaneously for all the considered time windows, with no adjustable parameters for the three smallest time windows [7].

We would also like to stress that, according to our theory, the value of the exponent $\mu \approx 1.6$ used in (1) to fit the tails of the distributions of seismic rates is describing a cross-over rather than a genuine asymptotic tail. Recall that the distribution of the total number of aftershocks has two power law regimes $\sim 1/r^{1+\frac{1}{\gamma}}$ for $r < r^* \simeq 1/(1-n)^{\gamma/(\gamma-1)}$ and $\sim 1/r^{1+\gamma}$ for $r > r^*$ [9]. The existence of this cross-over together with the concave shape of the distribution at small and intermediate values of r combine to create an effective power law with an apparent exponent $\mu \approx 1.6$ larger than the largest asymptotic exponent γ . We have verified this to be the case in synthetically generated distributions with genuine asymptotics exponent $\gamma = 1.25$ for instance, which could be well fitted by $\mu \approx 1.6$ over several decades. We note also that Pisarenko and Golubeva [11], with a different approach applied to much larger spatial box sizes in California, Japan and Pamir-Tien Shan, have reported an exponent $\mu < 1$ which could be associated with the intermediate asymptotics characterized by the exponent $1/\gamma < 1$, found in our previous analysis [10]. By using data collapse with varying spatial box sizes on a California catalog, Corral finds that the distribution of seismic rates exhibits a double power-law behavior with $\mu \approx 0$ for small rates and $\mu \approx 1.2$ for large rates [6]. The first regime might be associated with the non universal bulk part of the distribution found in our analysis. The second

regime is compatible with the prediction for the asymptotic exponent $\mu = \gamma$. In conclusion, we have offered a simple explanation of the power law distribution of seismic rates, which is derived from the other known power laws and the physics of cascades of earthquake triggering.

This work was partially supported by NSF-EAR02-30429 and by the Southern California Earthquake Center funded by NSF. —————

* Electronic address: sornette@moho.ess.ucla.edu

- [1] Helmstetter, A., Phys. Rev. Lett. 91, 058501 (2003).
- [2] Sornette, D. and P. Davy, Geophys. Res. Lett. 18, 1079 (1991).
- [3] Davy, P. et al., Nature 348, 56 (1990).
- [4] Kagan, Y.Y. and L. Knopoff, Geophys. J. Roy. Astr. Soc., 62, 303 (1980).
- [5] Kagan, Y.Y., Nonlin. Proc. Geophys., 1, 171, 1994.
- [6] Bak, P. et al., Phys. Rev. Lett. 88, 178501 (2002); Corral A., Phys. Rev. E. 6803(3 Part 2), 5102 (2003).
- [7] Saichev, A. and D. Sornette, physics/0412043
- [8] Helmstetter, A. and D. Sornette, J. Geophys. Res., 108 (B10), 2457 10.1029/2003JB002409 01, 2003.
- [9] Saichev, A. et al. cond-mat/0305007.
- [10] Saichev, A. and D. Sornette, Phys. Rev. E 70, 046123 (2004).
- [11] V.F. Pisarenko and T.V. Golubeva, Comp. Seism. Geodyn., 4, 127 (1996).